Initial U.S. Approval: 1999

WARNING: MYELOSUPPRESSION

See full prescribing information for complete boxed warning

- Causes severe and prolonged myelosuppression. (5.1)
- Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complications of the prolonged myelosuppression (5.1)

- INDICATIONS AND USAGE

Busulfan injection is an alkylating drug indicated for:

Use in combination with cyclophosphamide as a conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation for chronic myelogenous leukemia (CML) (1)

- DOSAGE AND ADMINISTRATION -

- Pre-medicate with anticonvulsants (e.g. benzodiazepines, phenytoin, valproic acid or levetracetam) and antiemetic (2.1, 5.2)

 Dilute and administer as intravenous infusion. Do not administer as intravenous push or
- bolus (2.1, 2.3)
- Recommended adult dose: 0.8 mg per kg of ideal body weight or actual body weight, whichever is lower, administered intravenously via a central venous catheter as a two-hour infusion every six hours for four consecutive days for a total of 16 doses (2.1)

– DOSAGE FORMS AND STRENGTHS

• 60 mg per 10 mL (6 mg per mL) single-dose vial (3)

FULL PRESCRIBING INFORMATION: CONTENTS* WARNING: MYELOSUPPRESSION

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
- Initial Dosing Information
- Preparation and Administration Precautions Preparation for Intravenous Administration
- 3. DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
- Myelosuppression Seizures
- Hepatic Veno-Occlusive Disease (HVOD)
- Embryo-fetal Toxicity
- Cardiac Tamponade Bronchopulmonary Dysplasia
- 5.7 Cellular Dysplasia
- 6. ADVERSE REACTIONS

- 6.1 Clinical Trial Experience
 6.2 Postmarketing Experiences
 6.3 Oral Busulfan Literature Review
- DRUG INTERACTIONS

60 mg/10 mL

4500415

- 7.1 Drugs that Decrease Busulfan Clearance 7.2 Drugs that Increase Busulfan Clearance

FULL PRESCRIBING INFORMATION

WARNING: MYLOSUPPRESSION Busulfan Injection causes severe and prolonged myelosuppression at the recommended dosa Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complicati of the prolonged myelosuppression. [See Warnings and Precautions (5.1)].

INDICATIONS AND USAGE

Busulfan injection is indicated for use in combination with cyclophosphamide as a conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation for chronic myeloge-

2. DOSAGE AND ADMINISTRATION

- Initial Dosing Information
 Administer busulfan injection in combination with cyclophosphamide as a conditioning regimen prior to bone marrow or peripheral blood progenitor cell replacement. For patients weighing more than 12 kg, the recommended doses are:
 - ghing more than 12 kg, the recommended doses are:
 Busulfan injection 0.8 mg per kg (ideal body weight or actual body weight, whichever is lower) intravenously via a central venous catheter as a two-hour infusion every six hours for four consecutive days for a total of 16 doses (Days -7, -6, -5 and -4).
 Cyclophosphamide 60 mg per kg intravenously as a one-hour infusion on each of two days beginning no sooner than six hours following the 16th dose of busulfan injection (Days -3 and -2).

 Administrate hematopoiatic progenitor cells on Day 0.

 - Administer hematopoietic progenitor cells on Day 0.
- Premedicate patients with anticonvulsants (e.g., benzodiazepines, phenytoin, valproic acid or levetiracetam) to prevent seizures reported with the use of high dose busulfan injection. Administer anticonvulsants 12 hours prior to busulfan injection to 24 hours after the last dose of busulfan injection [see Warnings and Precautions (5.2)].
- Administer antiemetics prior to the first dose of busulfan injection and continue on a fixed
- Schedule through busulfan injection administration.

 Busulfan injection clearance is best predicted when the busulfan injection dose is administration. tered based on adjusted ideal body weight. Dosing busulfan injection based on actual body weight, ideal body weight or other factors can produce significant differences in busulfan injection clearance among lean, normal and obese patients.
- o Calculate ideal body weight (IBW) as follows (height in cm, and weight in kg):
 Men: IBW (kg)=50+0.91x (height in cm -152)
 Women: IBW (kg)=45+0.91x (height in cm -152)
- For obese or severely obese patients, base busulfan injection dosing on adjusted ideal body weight (AIBW):

 AIBW=IBW +0.25x (actual weight -IBW).

2.2 Preparation and Administration PrecautionsBusulfan injection is incompatible with polycarbonate. Do not use any infusion components (syringes, filter needles, intravenous tubing, etc.) containing polycarbonate with busulfan injection. Use an administration set with minimal residual hold-up volume (2mL - 5mL) for product

Busulfan injection is a cytotoxic drug. Follow applicable special handling and disposal procedures. Skin reactions may occur with accidental exposure. Use gloves when preparing busulfan injection. If busulfan or diluted busulfan solution contacts the skin or mucosa, wash the skin or mucosa thoroughly with water.

Visually inspect parenteral drug products for particulate matter and discoloration prior to administration whenever the solution and container permit. Do not use if particulate matter is seen in the busulfan injection vial.

2.3 Preparation for Intravenous Administration

(56 mg total dose).

2.3 Preparation for intravenous Administration
Busulfan injection must be diluted prior to intravenous infusion with either 0.9% Sodium Chloride
Injection, USP (normal saline) or 5% Dextrose Injection, USP (DSW). The diluent quantity should
be 10 times the volume of busulfan injection, so that the final concentration of busulfan is
approximately 0.5 mg per mL. Calculation of the dose for a 70 kg patient would be performed

(70 kg patient) x (0.8 mg per kg) ÷ (6 mg per mL) =9.3 mL busulfan injection

To prepare the final solution for infusion, add 9.3 mL of busulfan injection to 93 mL of diluent (normal saline or DSW) as calculated below:

(9.3 mL busulfan injection) x (10) =93 mL of either diluent plus the 9.3 mL of

busulfan injection to yield a final concentration of busulfan of 0.54 mg per mL $(9.3 \text{ mL x 6 mg per mL} \div 102.3 \text{ mL} = 0.54 \text{ mg per mL}).$

- CONTRAINDICATIONS -

Busulfan injection is contraindicated in patients with a history of hypersensitivity to any of its components (4)

- WARNINGS AND PRECAUTIONS -Seizures: Initiate anticonvulsant prophylactic therapy prior to treatment with busulfan injection. Monitor patients with history of seizure disorder, head trauma or receiving
- replieptogenic drugs (5.2)
 Hepatic Veno-Occlusive Disease (HVOD): Increased risk of developing HVOD at AUC greater than 1,500 µM·min. Monitor serum transaminases, alkaline phosphatase and bilirubin daily (5.3)
- Embryo-fetal Toxicity: Can cause fetal harm. Advise of potential risk to a fetus and use of effective contraception (5.4, 8.1, 8.3)
- Cardiac tamponade has been reported in pediatric patients with thalassemia who received high doses of oral busulfan and cyclophosphamide. Abdominal pain and vomiting precede the tamponade in most patients (5.5)

ADVERSE REACTIONS —

Most common adverse reactions (incidence > 60%) were: myelosuppression, nausea, stomatitis, vomiting, anorexia, diarrhea, insomnia, fever, hypomagnesemia, abdominal pain, anxiety, headache, hyperglycemia and hypokalemia (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. --- DRUG INTERACTIONS -

- Drugs that Decrease busulfan Clearance: Metronidazole, itraconazole, Iron chelating agent, acetaminophen. (7.1)
 Drugs that Increases busulfan Clearance: Phenytoin. (7.2)
- ---- USE IN SPECIFIC POPULATIONS -
- Lactation: Discontinue breastfeeding (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

8. USE IN SPECIFIC POPULATIONS

- Pregnancy Lactation
 - Females and Males of Reproductive Potential Pediatric Use
- 8.5 Geriatric Use
- 10. OVERDOSAGE
- 11. DESCRIPTION
- 12. CLINICAL PHARMACOLOGY
 - 12.3 Pharmacokinetics
- 13. NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14. CLINICAL STUDIES
- 15. REFERENCES
- HOW SUPPLIED/STORAGE AND HANDLING
- 16.2 Storage and Handling
- 17. PATIENT COUNSELING INFORMATION

Sections or subsections omitted from the full prescribing information are not listed

All transfer procedures require strict adherence to aseptic techniques, preferably employing a vertical laminar flow safety hood while wearing gloves and protective clothing.

DO NOT put the busulfan injection into an intravenous bag or large-volume syringe that does not contain normal saline or D5W. Always add the busulfan injection to the diluent, not the diluent to the busulfan injection. Mix thoroughly by inverting several times.

Infusion pumps should be used to administer the diluted busulfan injection solution. Set the flow rate of the pump to deliver the entire prescribed busulfan injection dose over two hours. Prior to and following each infusion, flush the indwelling catheter line with approximately 5 mL of 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection, USP. DO NOT infuse concomitantly with another intravenous solution of unknown compatibility. WARNING: RAPID INFUSION OF BUSULFAN INJECTION HAS NOT BEEN TESTED AND IS NOT RECOMMENDED.

DOSAGE FORMS AND STRENGTHS

Busulfan Injection is supplied as a clear, colorless, sterile, solution in 10 mL single-dose vial containing 60 mg of busulfan at a concentration of 6 mg per mL for *intravenous use only*.

CONTRAINDICATIONS

Busulfan is contraindicated in patients with a history of hypersensitivity to any of its components

WARNINGS AND PRECAUTIONS

5.1 Myelosuppression

The most frequent serious consequence of treatment with busulfan at the recommended dos and schedule is prolonged myelosuppression, occurring in all patients (100%). Severe granulo-cytopenia, thrombocytopenia, anemia, or any combination thereof may develop. Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complications of the prolonged myelosuppression. Monitor complete blood counts, including white blood cell differentials, and quantitative platelet counts daily during treatment and until engraftment is demonstrated. Absolute neutrophil counts dropped below 0.5x10°/L at a median of 4 days post-transplant in 100% of patients treated in the busulfan clinical trial. The absolute neutrophil count recovered at a median of 13 days following allogeneic transplantation when prophylactic filgrastim was used in the majority of patients. Thrombocytopenia (less than 25,000/mm³ or requiring platelet transfusion) occurred at a median of 5-6 days in 98% of patients. Anemia (hemoglobin less than 8.0 g/dL) occurred in 69% of patients. Use antibiotic therapy and platelet and red blood cell

Seizures have been reported in patients receiving high-dose oral busulfan at doses producing plasma drug levels similar to those achieved following the recommended dosage of plasma drug levels similar to those achieved following the recommended dosage of busulfan. Despite prophylactic therapy with phenytoin, one seizure (1/42 patients) was reported during an autologous transplantation clinical trial of busulfan. This episode occurred during the cyclophosphamide portion of the conditioning regimen, 36 hours after the last busulfan dose. Initiate phenytoin therapy or any other alternative anti-convulsant prophylactic therapy (e.g. benzodiazepines, valproic acid or levetiracetam) prior to busulfan treatment (See Dosage and Administration (2.1)]. Use caution when administering the recommended dose of busulfan to patients with a history of a seizure disorder or head trauma or who are receiving other potentially patients with a history of a seizure disorder or head trauma or who are receiving other potentially epileptogenic drugs.

5.3 Hepatic Veno-Occlusive Disease (HVOD)

Current literature suggests that high busulfan area under the plasma concentration verses time curve (AUC) values (greater than 1,500 µM·min) may be associated with an increased risk of developing HVOD. Patients who have received prior radiation therapy, greater than or equal to three cycles of chemotherapy, or a prior progenitor cell transplant may be at an increased risk of developing HVOD with the recommended busulfan dose and regimen. Based on clinical examination and laboratory findings, HVOD was diagnosed in 8% (5/61) of patients treated with busulfan in the setting of allogeneic transplantation, was fatal in 2/5 cases (40%), and yielded an overall mortality from HVOD in the entire study population of 2/61 (3%). Three of the five patients diagnosed with HVOD were retrospectively found to meet the Jones' criteria. The incidence of HVOD reported in the literature from the randomized, controlled trials was 7.7%-12% [See Clinical Studies (14)1. Monitor serum transaminases, alkaline phosphatase, and bilirubin daily through BMT Day +28 to detect hepatotoxicity, which may herald the onset of HVOD.

5.4 Embryo-fetal Toxicity

Busulfan can cause fetal harm when administered to a pregnant woman based on animal data.

Busulfan was teratogenic in mice, rats, and rabbits. The solvent, DMA, may also cause fetal harm when administered to a pregnant woman based on findings in animals. Advise pregnant woman based on findings in animals. Advise pregnant woman based on pregnant woman programment of percentilizing the affairs. Advise formulas and males of percentilizing the pregnant woman based on programment of the pregnant woman based on programment of the progra women of the potential risk to a fetus. Advise females and males of reproductive potential to use effective contraception during and after treatment with busulfan [see Use in Specific Populations

5.5 Cardiac Tamponade

Cardiac tamponade has been reported in pediatric patients with thalassemia (8/400 or 2% in one series) who received high doses of oral busulfan and cyclophosphamide as the preparatory regimen for hematopoietic progenitor cell transplantation. Six of the eight children died and two were saved by rapid pericardiocentesis. Abdominal pain and vomiting preceded the tamponade n most patients. Monitor for signs and symptoms, promptly evaluate and treat if cardiac

5.6 Bronchopulmonary DysplasiaBronchopulmonary dysplasia with pulmonary fibrosis is a rare but serious complication following chronic busulfan therapy. The average onset of symptoms is 4 years after therapy (range 4 months to 10 years).

5.7 Cellular Dysplasia

Busulfan may cause cellular dysplasia in many organs. Cytologic abnormalities characterized by giant, hyperchromatic nuclei have been reported in lymph nodes, pancreas, thyroid, adrenal glands, liver, lungs and bone marrow. This cytologic dysplasia may be severe enough to cause difficulty in the interpretation of exfoliative cytologic examinations of the lungs, bladder, breast and the uterine cervix.

ADVERSE REACTIONS

e following adverse reactions are discussed in more detail in other sections of the labeling: Myelosuppression [See Warnings and Precautions (5.1)] Seizures [See Warnings and Precautions (5.2)]

- Hepatic Veno-Occlusive Disease (HVOD) [See Warnings and Precautions (5.3)] Embryo-fetal Toxicity [See Warnings and Precautions (5.4)] Cardiac Tamponade [See Warnings and Precautions (5.5)]
- Bronchopulmonary Dysplasia [see Warnings and Precautions (5.6)] Cellular Dysplasia [See Warnings and Precautions (5.7)]

6.1 Clinical Trial Experience

Non-Hematological Adverse Reactions

BODY AS A WHOLE

Revised: 06/2019

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reaction information is primarily derived from the clinical study (N=61) of busulfan and the data obtained for high-dose oral busulfan conditioning in the setting of randomized, controlled trials identified through a literature review.

In the busulfan Injection allogeneic stem cell transplantation clinical trial, all patients were treated with busulfan 0.8 mg per kg as a two-hour infusion every six hours for 16 doses over four days, combined with cyclophosphamide 60 mg per kg x2 days. Ninety-three percent (93%) of evaluable patients receiving this dose of busulfan maintained an AUC less than 1,500 µM·min for dose 9, which has generally been considered the level that minimizes the risk of HVOD.

Table 1 lists the non-hematologic adverse reactions events through Bone Marrow Transplantation (BMT) Day +28 at a rate greater than or equal to 20% in patients treated with busulfan prior to allogeneic hematopoietic cell transplantation.

Table 1: Summary of the Incidence (greater than or equal to 20%) of Non-Hematologic Adverse Reactions through BMT Day +28 in Patients who Received Busulfan injection Prior to Allogeneic Hematopoietic Progenitor Cell Transplantation

Percent Incidence

DOD I AO A WITOLL	
Fever	80
Headache	69
Asthenia	51
Chills	46
Pain	44
Edema General	28
Allergic Reaction	26
Chest Pain	26
Inflammation at Injection Site	25
Back Pain	23
CARDIOVASCULAR SYSTEM	
Tachycardia	44
Hypertension	36
Thrombosis	33
Vasodilation	25
DIGESTIVE SYSTEM	
Nausea	98
Stomatitis (Mucositis)	97
Vomiting	95
Anorexia	85
Diarrhea	84
Abdominal Pain	72
Dyspepsia	44
Constipation	38
Dry Mouth	26
Rectal Disorder	25
Abdominal Enlargement	23
METABOLIC AND NUTRITIONAL SYSTEM	20
Hypomagnesemia	77
Hyperglycemia	66
Hypokalemia	64
Hypocalcemia	49
Hyperbilirubinemia	49
Edema	36
SGPT Elevation	30
	•
Creatinine Increased NERVOUS SYSTEM	21
	0.4
Insomnia	84
Anxiety	72
Dizziness	30
Depression	23
RESPIRATORY SYSTEM	
Rhinitis	44
Lung Disorder	34
Cough	28
Epistaxis	25
Dyspnea	25
SKIN AND APPENDAGES	
Rash	57
Pruritus	28
1. Includes all reported adverse reactions regardles	ss of severity (toxicity grades 1-4)

Additional Adverse Reactions by Body System

Hematologic: Prolonged prothrombin time

Gastrointestinal: Esophagitis, ileus, hematemesis, pancreatitis, rectal discomfort

Hepatic: Alkaline phosphatase increases, jaundice, hepatomegaly Graft-versus-host disease: Graft-versus-host disease. There were 3 deaths (5%) attributed

Edema: Hypervolemia, or documented weight increase

Infection: Infection, pneumonia (fatal in one patient and life-threatening in 3% of patients)

Cardiovascular: Arrhythmia, atrial fibrillation, ventricular extrasystoles, third degree heart block, thrombosis (all episodes were associated with the central venous catheter), hypotension. flushing and hot flashes, cardiomegaly, ECG abnormality, left-sided heart failure, and pericardial

Pulmonary: Hyperventilation, alveolar hemorrhage (fatal in 3%), pharyngitis, hiccup, asthma, atelectasis, pleural effusion, hypoxia, hemoptysis, sinusitis, and interstitial fibrosis (fatal in a single case)

Neurologic: Cerebral hemorrhage, coma, delirium, agitation, encephalopathy, confusion, hallucinations, lethargy, somnolence

Renal: BUN increased, dysuria, oliguria, hematuria, hemorrhagic cystitis

Skin: Alopecia, vesicular rash, maculopapular rash, vesiculo-bullous rash, exfoliative dermatitis, erythema nodosum, acne, skin discoloration

6.2 Postmarketing ExperienceBecause these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The following adverse reactions have been identified during post-approval use of Purulfan lineties: Busulfan Injection:

Blood and Lymphatic System Disorders: febrile neutropenia

Gastrointestinal Disorders: tooth hypoplasia

Metabolism and Nutrition Disorders: tumor lysis syndrome

Vascular Disorders: thrombotic microangiopathy (TMA)

Infections and Infestations: severe bacterial, viral (e.g., cytomegalovirus viremia) and fungal infections; and sepsis.

6.3 Oral Busulfan Literature Review

A literature review identified four randomized, controlled trials that evaluated a high-dose oral busulfan-containing conditioning regimen for allogeneic bone marrow transplantation in the setting of CML *[see Clinical Studies (14)]*. The safety outcomes reported in those trials are summarized in Table 2 below for a mixed population of hematological malignancies (AML, CML,

Table 2: Summary of safety analyses from the randomized, controlled trials utilizing a high dose oral busulfan-containing conditioning regimen that were identified in a literature review.

		(Clift		
		CML Chr	onic Phase		
TRM ¹	VOD ²	GVHD ³	Pulmonary	Hemorrhagic Cystitis	Seizure
Death ≤100d =4.1% (3/73)	No Report	Acute ≥Grade 2 =35% Chron- ic=41% (30/73)	1 death from Idiopathic Interstitial Pneumonitis And 1 death from Pulmonary Fibrosis	No Report	No Report
		Dev	vergie		
		CML Chr	onic Phase		
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
38%	7.7% (5/65) Deaths=4.6% (3/65)	Acute ≥Grade 2 =41% (24/59 at risk)	Interstitial Pneumonitis= 16.9% (11/65)	10.8% (7/65)	No Report
		Rin	ngden		
		CML, A	AML, ALL		
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
28%	12%	Acute ≥Grade 2 GVHD=26% Chronic GVHD =45%	Interstitial Pneumonitis =14%	24%	6%
		BI	lume		
		CML,	AML,ALL		
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
No Report	Deaths =4.9%	Acute ≥Grade 2 GVHD=22% (13/58 at risk) Chronic GVHD =31% (14/45 at risk)	No Report	No Report	No Report

7.1 Drugs that Decrease Busulfan Clearance Itraconazole decreases busulfan clearance by up to 25%. Metronidazole decreases the clearance of busulfan to a greater extent than does itraconazole; metronidazole coadministration has been associated with increased busulfan toxicity. Fluconazole (200 mg) has been used

mechanism of this interaction is not fully elucidated. Discontinue iron chelating agents well in advance of administration of busulfan to avoid increased exposure to busulfan.

Busulfan can cause fetal harm when administered to a pregnant woman based on animal data. Busulfan was teratogenic in mice, rats, and rabbits following administration during organogenesis. The solvent, DMA, may also cause fetal harm when administered to a pregnant woman. In rats, DMA doses of approximately 40% of the daily dose of DMA in the busulfan dose on a mg/m² basis given during organogenesis caused significant developmental anomalies [see Data]. There are no available human data informing the drug-associated risk. Advise pregnant

7. DRUG INTERACTIONS

Because busulfan is eliminated from the body via conjugation with glutathione, use of acetamin-ophen prior to (less than 72 hours) or concurrent with busulfan may result in reduced busulfan clearance based upon the known property of acetaminophen to decrease glutathione levels in

of glutathione-S-transferase. Since the pharmacokinetics of busulfan was studied in patients or gutationic variation and the clearance of busulfan at the recommended dose may be lower and exposure (AUC) higher in patients not treated with phenytoin.

8.1 Pregnancy Risk Summary

women of the potential risk to a fetus.

Following administration during organogenesis in animals, busulfan caused malformations and anomalies, including significant alterations in the musculoskeletal system, body weight gain,

2. VOD = Veno-Occlusive Diseases of the Liver 3. GVHD = Graft versus Host Disease

Decreased clearance of busulfan was observed with concomitant use with deferasirox. The

7.2 Drugs that Increase Busulfan ClearancePhenytoin increases the clearance of busulfan by 15% or more, possibly due to the induction

8. USE IN SPECIFIC POPULATIONS

The background risk of major birth defects and miscarriage for the indicated populations are unknown. However, the background risk in the U.S. general population of major birth defects is 2-4% and of miscarriage is 15-20% of clinically recognized pregnancies.

Metabolic: Hypophosphatemia, hyponatremia

Other Events: Injection site pain, myalgia, arthralgia, ear disorder

and size. In pregnant rats, busulfan produced sterility in both male and female offspring due to the absence of germinal cells in the testes and ovaries. The solvent, N,N-dimethylacetamide (DMA), administered to rats at doses of 400 mg/kg/day (about 40% of the daily dose of DMA in the busulfan dose on a mg/m² basis) during organogenesis caused significant developmental anomalies. The most striking abnormalities included anasarca, cleft palate, vertebral anomalies, rib anomalies, and serious anomalies of the vessels of the heart.

8.2 Lactation

It is not known whether busulfan is present in human milk. Because many drugs are excreted in human milk and because of the potential for tumorigenicity shown for busulfan in human and animal studies, discontinue breastfeeding during treatment with busulfan

8.3 Females and Males of Reproductive Potential

Females
Busulfan can cause fetal harm when administered to a pregnant woman [see Use in Specific Populations (8.1)1. Advise females of reproductive potential to use effective contraception during treatment with busulfan and for 6 months following cessation of therapy

Males
Busulfan may damage spermatozoa and testicular tissue, resulting in possible genetic fetal abnormalities. Males with female sexual partners of reproductive potential should use effective contraception during treatment with busulfan and for 3 months after cessation of therapy [see

Infertility

<u>Females</u>
Ovarian suppression and amenorrhea commonly occur in premenopausal women undergoing orania suppression and amendment commonly occur in preintendpassal women undergoned chronic, low-dose busulfan therapy for chronic myelogenous leukemia. Busulfan may cause temporary or permanent infertility in prepubertal girls or in females of child-bearing potential treated with high-dose busulfan in the conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation

Males Sterility, azoospermia, and testicular atrophy have been reported in male patients.

8.4 Pediatric Use

The effectiveness of busulfan in the treatment of CML has not been specifically studied in pediatric patients. An open-label, uncontrolled study evaluated the pharmacokinetics of busulfar in 24 pediatric patients receiving busulfan as part of a conditioning regimen administered prior to hematopoietic progenitor cell transplantation for a variety of malignant hematologic (N=15 or non-malignant diseases (N=9). Patients ranged in age from 5 months to 16 years (median 3 years). Busulfan dosing was targeted to achieve an area under the plasma concentration curve (AUC) of 900-1350 µ.M·min with an initial dose of 0.8 mg per kg or 1.0 mg per kg (based on Actual Body Weight (ABW)) if the patient was greater than 4 or less than or equal to 4 years, respectively. The dose was adjusted based on plasma concentration after completion of dose 1

Patients received busulfan doses every six hours as a two-hour infusion over four days for a rations received obstillar to obses every six hours as a two-floor influsion over four days for total of 16 doses, followed by cyclophosphamide 50 mg per kg once daily for four days. After one rest day, hematopoietic progenitor cells were inflused. All patients received phenytoin as seizure prophylaxis. The target AUC (900-1350±5% μ.M-min) for busulfan was achieved at dose 1 in 71% (17/24) of patients. Steady state pharmacokinetic testing was performed at dose 9 and 13. Busulfan levels were within the target range for 21 of 23 evaluable patients.

All 24 patients experienced neutropenia (absolute neutrophil count (ANC) less than 0.5x109/L) All 24 patients experienced neutropenia (absolute neutrophii count (ANC) less man 0.5x10^{-y}l and thrombocytopenia (platelet transfusions or platelet count less than 20,000/mm³). Seventy-nine percent (19/24) of patients experienced lymphopenia (absolute lymphocyte count less than 0.1x10⁹). In 23 patients, the ANC recovered to greater than 0.5x10⁹/L (median time to recovery = BMT day +13; range = BMT day +9 to +22). One patient who died on day +20 had not recovered to an ANC >0.5x109/I

Four (17%) patients died during the study. Two patients died within 28 days of transplant; one with pneumonia and capillary leak syndrome, and the other with pneumonia and veno-occlusive disease. Two patients died prior to day 100; one due to progressive disease and one due to

Adverse reactions were reported in all 24 patients during the study period (BMT day -10 through BMT day +28) or post-study surveillance period (day +29 through +100). These included vomiting (100%), nausea (83%), stomatitis (79%), HVOD (21%), graft-versus host disease (GVHD) (25%), and pneumonia (21%).

Based on the results of this 24-patient clinical trial, a suggested dosing regimen of busulfan ir pediatric patients is shown in the following dosing nomogram:

Busulfan Dosing Nomogram				
Patient's Actual Body Weight (ABW)	Busulfan Dosage			
less than or equal to12 kg	1.1 (mg per kg)			
greater than 12 kg	0.8 (mg per kg)			

Simulations based on a pediatric population pharmacokinetic model indicate that approximately 60% of pediatric patients will achieve a target busulfan exposure (AUC) between 900 to 1350 u.M. min with the first dose of busulfan using this dosing nomogram. The appeutic drug monitoring and dose adjustment following the first dose of busulfan is recommended

Dose Adjustment Based on Therapeutic Drug Monitoring
Instructions for measuring the AUC of busulfan at dose 1 (see Blood Sample Collection for AUC Determination) and the formula for adjustment of subsequent doses to achieve the desired target AUC (1125 µM•min), are provided below.

Adjusted dose (mg) = Actual Dose (mg) x Target AUC (μM•min)/Actual AUC (μΜ•min) For example, if a patient received a dose of 11 mg busulfan and if the corresponding AUC measured was $800~\mu\text{M} \cdot \text{min}$, for a target AUC of 1125 $\mu\text{M} \cdot \text{min}$, the target mg dose would be:

Mg dose = 11 mg x 1125 μ M•min /800 μ M•min = 15.5 mg Busulfan dose adjustment may be made using this formula and instructions below.

Blood Sample Collection for AUC Determination

Calculate the AUC (μM•min) based on blood samples collected at the following time points:

For dose 1:2 hr (end of infusion), 4 hr and 6 hr (immediately prior to the next scheduled busulfan administration). Actual sampling times should be recorded.

For doses other than dose 1: Pre-infusion (baseline), 2 hr (end of infusion), 4 hr and 6 hr tely prior to the next scheduled busulfan administra

AUC calculations based on fewer than the three specified samples may result in inaccurate

AUC determinations. For each scheduled blood sample, collect one to three mL of blood into heparinized (Na or Li heparin) Vacutainer® tubes. The blood samples should be placed on wet ice immediately after collection and should be centrifuged (at 4°C) within one hour. The plasma, harvested into appropriate cryovial storage tubes, is to be frozen immediately at -20°C. All plasma samples are to be sent in a frozen state (i.e., on dry ice) to the assay laboratory for the determination of

Calculation of AUC

Busulfan AUC calculations may be made using the following instructions and appropriate standard pharmacokinetic formula

Dose 1 AUC $_{infinity}$ Calculation: AUC $_{infinity}$ = AUC $_{o.6hr}$ + AUC $_{extrapolated}$, where AUC $_{o.6hr}$ is to be estimated using the linear trapezoidal rule and AUC extrapolated can be computed by taking the ratio of the busulfan concentration at Hour 6 and the terminal elimination rate constant, λ_z . The λ_z must be calculated from the terminal elimination phase of the busulfan concentration vs. time curve. A "0" pre-dose busulfan concentration should be assumed, and used in the calculation of AUC.

If the AUC is assessed subsequent to Dose 1, steady-state AUC_{ss} (AUC_{0-6hr}) is to be estimated from the trough, 2 hr, 4 hr and 6 hr concentrations using the linear trapezoidal rule Instructions for Drug Administration and Blood Sample Collection for Therapeutic Drug

Use an administration set with minimal residual hold up (priming) volume (1 to 3 mL) for drug infusion to ensure accurate delivery of the entire prescribed dose and to ensure accurate collec-

tion of blood samples for therapeutic drug monitoring and dose adjustment Prime the administration set tubing with drug solution to allow accurate documentation of the

start time of busulfan infusion. Collect the blood sample from a peripheral IV line to avoid

contamination with infusing drug. If the blood sample is taken directly from the existing central venous catheter (CVC), <u>DO NOT COLLECT THE BLOOD SAMPLE WHILE THE DRUG IS</u> INFUSING to ensure that the end of infusion sample is not contaminated with any residual drug. At the end of infusion (2 hr), disconnect the administration tubing and flush the CVC line with At the end of influsion (2 m), assorbled the administration tubing and lists in each collection of the end of influsion sample from the CVC por Collect the blood samples from a different port than that used for the busulfan infusion. Whe recording the busulfan infusion stop time, do not include the time required to flush the indwelling catheter line. Discard the administration tubing at the end of the two-hour infusion [see Dosage

Clinical studies of busulfan did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects

10 OVERDOSAGE

here is no known antidote to busulfan injection other than hematopoietic progenitor cell transplantation. In the absence of hematopoietic progenitor cell transplantation, the recommended dosage for busulfan would constitute an overdose of busulfan. The principal toxic effect is ubasage for businaria would constitute an overdose or businian. The principal roots effect is profound bone marrow hypoplasia/aplasia and pancytopenia, but the central nervous system liver, lungs, and gastrointestinal tract may be affected. Monitor hematologic status closely and institute vigorous supportive measures as medically indicated. Survival after a single 140 mg dose of Myleran® Tablets in an 18 kg, 4-year old child has been reported. Inadvertent administra-tion of a greater than normal dose of oral busulfan (2.1 mg per kg) total dose of 23.3 mg per kg) occurred in a 2-year old child prior to a scheduled bone marrow transplant without sequelae. An acute dose of 2.4 g was fatal in a 10-year old boy. There is one report that busulfan is dialyzable, thus dialysis should be considered in the case of overdose

11 DESCRIPTION

Ifan is a bifunctional alkylating agent known chemically as 1,4-butanediol, dimethanesul fonate. Busulfan Injection is intended for intravenous administration. It is supplied as a clear. colorless, sterile, solution in 10 mL single-dose vials. Each vial of busulfan injection contains 60 mg (6 mg/mL) of busulfan, the active ingredient, a white crystalline powder with a molecular formula of CH₃SO₂O(CH₂)₄OSO₂CH₃ and a molecular weight of 246 g/mole. Busulfan has the following chemical structure:

$$CH_3 - S - O - CH_2 - CH_2 - CH_2 - CH_2 - O - S - CH_3$$

Rusulfan is dissolved in N N-dimethylacetamide (DMA), 3.3 ml, and Polyethylene Glycol 400. businant's dissolved in Ny-dimetrifaceating (billy), 3.5 file. and Polyetrifete Glycul 40th NF 6.7 mL. The solubility of busulfan in water is 0.1 g per L and the pH of busulfan diluted to approximately 0.5 mg per mL busulfan in 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection, USP as recommended for infusion reflects the pH of the diluent used and ranges

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.1 Mechanism of Action

Busulfan is a bifunctional alkylating agent in which two labile methanesulfonate groups are attached to opposite ends of a four-carbon alkyl chain. In aqueous media, busulfan hydrolyzes to release the methanesulfonate groups. This produces reactive carbonium ions that can alkylate DNA. DNA damage is thought to be responsible for much of the cytotoxicity of busulfan.

12.3 Pharmacokinetics

The pharmacokinetics of husulfan was studied in 59 natients participating in a prospective trial of a busulfan cyclophosphamide preparatory regimen prior to allogeneic hematopoietic progenitor stem cell transplantation. Patients received 0.8 mg/kg busulfan every six hours, for a total of 16 doses over four days. Fifty-five of fifty-nine patients (93%) administered busulfan maintaine AUC values below the target value (less than 1500 µM•min)

Table 3: Steady State Pharmacokinetic Parameters Following Busulfan

(0.8 mg per kg; N=59)						
	Mean	CV (%)	Range			
C _{max} (ng per mL)	1222	18	496-1684			
AUC (μM•min)	1167	20	556-1673			
CL (mL per min per kg) ¹ 2.52 25 1.49-4.31						
¹ Clearance normalized to actual body weight for all patients.						

Busulfan pharmacokinetics showed consistency between dose 9 and dose 13 as demonstrated by reproducibility of steady state C_{max} and a low coefficient of variation for this parameter

Distribution: Busulfan achieves concentrations in the cerebrospinal fluid approximately equal to those in plasma. Busulfan primarily binds to albumin (Mean \pm standard deviation=32.4 \pm 2.2%) Metabolism: Busulfan is predominantly metabolized by conjugation with glutathione, both spontaneously and by glutathione S-transferase (GST) catalysis. This conjugate undergoes exten-

Excretion: Following administration of 14C-labeled busulfan to humans, approximately 30% of the radioactivity was excreted into the urine over 48 hours; negligible amounts were recovered

Specific Populations

Pediatric Patients: In a pharmacokinetic study of busulfan injection in 24 pediatric patients, the population pharmacokinetic (PPK) estimates of busulfan injection for clearance (CL) and volume of distribution (V) were determined. For actual body weight, PPK estimates of CL and V were 4.04 L/hr per 20 kg (3.37 mL per min per kg; interpatient variability 23%); and 12.8 L per 20 kg (0.64 L per kg; interpatient variability 11%).

13. NONCLINICAL TOXICOLOGY

sive oxidative metabolism in the liver

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Busulfan is a mutagen and a clastogen. In in vitro tests it caused mutations in Salmonella typhimurium and Drosophila melanogaster. Chromosomal aberrations induced by busulfan have been reported in vivo (rats, mice, hamsters, and humans) and in vitro (rodent and human cells). The intravenous administration of busulfan (48 mg/kg given as biweekly doses of 12 mg/kg, or 30% of the total busulfan dose on a mg/m² basis) has been shown to increase the incide

Busulfan depleted oocytes of female rats and induced sterility in male rats and hamsters. The solvent DMA may also impair fertility. A DMA daily dose of 0.45 g/kg/day given to rats for nine days (equivalent to 44% of the daily dose of DMA contained in the recommended dose of busulfan on a mg/m² basis) significantly decreased spermatogenesis in rats. A single subcutaneous dose of 2.2 g/kg (27% of the total DMA dose contained in busulfan on a mg/m² basis) four days after inse mination terminated pregnancy in 100% of tested hamsters [see Use in Specific Populations (8.3)1

14 CLINICAL STUDIES

Documentation of the safety and efficacy of busulfan as a component of a conditioning regimen prior to allogeneic hematopoietic progenitor cell reconstitution is derived from two source

- analysis of a prospective clinical trial of busulfan that involved 61 patients diagnosed with various hematologic malignancies, and the published reports of randomized, controlled trials that employed high-dose oral busulfan
- as a component of a conditioning regimen for transplantation, which were identified in a literature review of five established commercial databases.

Prospective Clinical Trial of Busulfan: The prospective trial was a single-arm open-label study in 61 patients who received busulfan as part of a conditioning regimen for allogeneic hematopoietic stem cell transplantation. The study included patients with acute leukemia past first remission (first or subsequent relapse), with high-risk first remission, or with induction failure chronic myelogenous leukemia (CML) in chronic phase, accelerated phase, or blast crisis; primary refractory or resistant relapsed Hodgkin's disease or non-Hodgkin's lymphoma; and myelodysplastic syndrome. Forty-eight percent of patients (29/61) were heavily pretreated, defined as having at least one of the following: prior radiation, greater than or equal to 3 prior chemotherapeutic regimens, or prior hematopoietic stem cell transplant. Seventy-five percent of chemotherapeutic regimens, or prior hematopoietic stern patients (46/61) were transplanted with active disease.

Patients received 16 busulfan doses of 0.8 mg per kg every 6 hours as a two-hour infusion for 4 days, followed by cyclophosphamide 60 mg per kg once per day for two days (BuCy2 regimen). All patients received 100% of their scheduled busulfan regimen. No dose adjustments were made. After one rest day, allogeneic hematopoietic progenitor cells were infused. The efficacy parameters in this study were myeloablation (defined as one or more of the following: absolute neutrophil count [ANC] less than $0.5x10^{9}/L$, absolute lymphocytes count [ALC] less than $0.1x10^{9}/L$, thrombocytopenia defined as a platelet count less than $20.000/mm^{3}$ or a platelet transfusion requirement) and engraftment (ANC greater than or equal to $0.5x10^{9}/L$).

All nationts (61/61) experienced myeloablation. The median time to neutropenia was 4 days All evaluable patients (60/60) engrafted at a median of 13 days post-transplant (range 9 to 29 days); one patient was considered non-evaluable because he died of a fungal pneumonia 20 days after BMT and before engraftment occurred. All but 13 of the patients were treated with prophylactic G-CSF. Evidence of donor cell engraftment and chimerism was documented in all patients who had a chromosomal sex marker or leukemic marker (43/43), and no patient with chimeric evidence of allogeneic engraftment suffered a later loss of the allogeneic graft. There were no reports of graft failure in the overall study population. The median number of platelet transfusions per patient was 6, and the median number of red blood cell transfusions

Twenty-three patients (38%) relapsed at a median of 183 days post-transplant (range 36 to 406 days). Sixty-two percent of patients (38/61) were free from disease with a median follow-up of 269 days post-transplant (range 20 to 583 days). Forty-three patients (70%) were alive with r 209 days post-transplant (angle 20 to 30 days). Forty-time patients (10%) were anive with median follow up of 288 days post-transplant (range 51 to 583 days). There were two deaths efore BMT Day +28 and six additional patients died by BMT Day +100. Ten patients (16%) died after BMT Day +100, at a median of 199 days post-transplant (range 113 to 275 days).

Oral Busulfan Literature Review: Four publications of randomized, controlled trials that evaluated a high-dose oral busulfan-containing conditioning regimen (busulfan 4 mg/kg/d x4 days + cyclophosphamide 60 mg/kg/d x2 days) for allogeneic transplantation in the setting of CML were identified. Two of the studies (Clift and Devergie) had populations confined to CML in chronic phase that were randomized between conditioning with busulfan/cyclophosphamide (BU/CY) and cyclophosphamide/total body irradiation (CY/TBI). A total of 138 patients were treated with and cyclopnospharmidertotal body irradiation (CYTEI). A total of 136 patients were treated with BUI/CY in these studies. The populations of the two remaining studies (Ringden and Blume) included patients with CML, acute lymphoblastic leukemia (ALL), and acute myelogenous eukemia (AML). In the Nordic BMT Group study published by Ringden, et al. 57 patients had CML eurening (AML). Intervoluce with BU/CY. Patients with CML in chronic phase, accelerated and of those, 30 were treated with BU/CY. Patients with CML in chronic phase, accelerated phase, and blast crisis were eligible for this study. The participants with CML (34/122 patients) n a SWOG study published by Blume, et al., had disease beyond first chronic phase. Twenty of those CML patients were treated with BU/CY, and the TBI comparator arm utilized etoposide

Table 4 summarizes the efficacy analyses reported from these 4 studies.

Table 4: Summary of efficacy analyses from the randomized, controlled trials utilizing a Ifan-containing conditioning regimen identified in a literature review

		9					
			Clift, 1				
3 year Overall 3 year DFS Survival (p=0.43)				Relapse		Time to Engraftment (ANC greater than or equal to 500)	
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
80%	80%	71%	68%	13%	13%	22.6 days	22.3 days
Devergie, 1995 CML Chronic Phase;							
5 year Over (p=0			Relapse (Relative Risk analysis BU/CY: CY/TBI) (p=0.04)		Time to Engraftment (ANC greater than or equal to 500)		
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
60.6% ±11.7%	65.8% ±12.5%	59.1% ±11.8%	51.0% ±14%		10 .00-20.28)	None Given	None Given
Ringden, 1994 CML,AML,ALL;							
Surv	3 year Overall Survival Survival (p<0.03) 3 year Relapse Free Survival (p=0.065) Relapse (p=0.9)			Time to Er (ANC gre 50			
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
62%	76%	56%	67%	22%	26%	20 days	20 days
Blume, 1993¹ CML, AML, ALL; Relative Risk Analysis BU/CY: Etoposide/TBI							
RR of Mortality DFS				RR of Relapse (Relative Risk analysis		Time to Er	ngraftment

Blume, 1993¹ CML, AML, ALL; Relative Risk Analysis BU/CY: Etoposide/TBI							
RR of M		DF		RR of Relapse		Time to Engraftment	
BU/CY	Eto/TBI	BU/CY	Eto/TBI	BU/CY	Eto/TBI	BU/CY	Eto/TBI
	0.97% (95% CI=0.64-1.48)		Not Given		1.02 (95% CI=0.56-1.86)		Siven

- Eto=etoposide. TBI was combined with etoposide in the comparator arm of this study.
- CY = Cyclophosphamide
- TBI = Total Body Irradiation
- DFS = Disease Free Survival ANC = Absolute Neutrophil Count

15. REFERENCES

1. OSHA Hazardous Drugs. OSHA. [Accessed on June 18, 2014 from http://www.osha.gov/SLTC/hazardousdrugs/index.html

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Busulfan injection is packaged as a sterile solution in 10 mL single-dose clear glass vials each containing 60 mg of busulfan at a concentration of 6 mg per mL for intravenous use.

Busulfan injection is distributed as a unit of eight individually packed single-dose vials as follows:

Product Code	Unit of Sale	Strength	Each
261610	NDC 65219-160-10 Unit of 8 individual cartons.	60 mg per 10 mL (6 mg per mL)	NDC 65219-160-01 10 mL single-dose vials packed in individual cartons.

16.2 Storage and Handling

Unopened vials of busulfan injection must be stored under refrigerated conditions between 2°C to 8°C (36°F to 46°F)

Busulfan injection diluted in 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection USP is stable at room temperature (25°C) for up to 8 hours but the infusion must be completed.

Busulfan injection diluted in 0.9% Sodium Chloride Injection, USP is stable at refrigerated conditions (2°C to 8°C) for up to 12 hours but the infusion must be completed within that time. Busulfan injection is a cytotoxic drug. Follow applicable special handling and disposal proce-

17. PATIENT COUNSELING INFORMATION

Myelosuppression

Advise patients of the possibility of developing low blood cell counts and the need for hematopoi etic progenitor cell infusion. Instruct patients to immediately report to their healthcare provider fever develops [see Warnings and Precautions (5.1)].

Seizures

Advise patients of the possibility of seizures and that they will be given medication to prevent them. Patients should be asked to report a history of seizure or head trauma [see Warnings and Precautions (5.2)]. Hepatic Veno-Occlusive Disease (HVOD)

Advise patients of the risks associated with the use of busulfan injection as well as the plan for regular blood monitoring during therapy. Specifically inform patients of the following: The risk of veno-occlusive liver disease [see Warnings and Precautions (5.3)].

Embryo-fetal Toxicity Advise females of reproductive potential of the potential risk to a fetus and to inform their health care provider with a known or suspected pregnancy [see Warnings and Precautions (5.4) and Use in Specific Populations (8.1)].

Females of Reproductive Potential

Advise females of reproductive potential to use effective contraception during treatment with busulfan injection and for 6 months following cessation of therapy [see Use in Specific Popu-

Males of Reproductive Potential

in Specific Populations (8.2)1

Advise males with female sexual partners of reproductive potential to use effective contraception during treatment with busulfan injection and for 3 months following cessation of therapy [see Use in Specific Populations (8.3)1.

Lactation Advise females to discontinue breastfeeding during treatment with busulfan injection [see Use

Infertility Advise females and males of reproductive potential that busulfan injection may cause temporary or permanent infertility [see *Use in Specific Populations* (8.3)].

Cardiac Tamponade

Advise patients of the risk of cardiac tamponade. Instruct patients to report to their healthcare provider symptoms of abdominal pain and vomiting [see Warnings and Precautions (5.5)]

Bronchopulmonary Dysplasia

Advise patients of the possibility of bronchopulmonary dysplasia with pulmonary fibrosis with chronic busulfan injection therapy. Instruct patients to report symptoms of shortness of breath and cough to their healthcare provider. These symptoms could occur several months or years after therapy with busulfan injection [see *Warnings and Precautions* (5.6)].

Manufactured for



Lake Zurich, IL 60047 www.fresenius-kabi.com/us Made in India

451315

Issued: June 2019

5