
WARNING

Carboplatin Injection should be adminis tered under the supervision of a qualified physician experienced in the use of cancer chemotherapeutic agents. Appro priate management of therapy and complications is possible only when adequate reatment facilities are readily available. Bone marrow suppression is dose related and may be severe, resulting in infection and/or bleeding. Anemia may be cumulative and may require transfusion support. Vomiting is another frequent drug-related side effect Anaphylactic-like reactions to carboplatin have been reported and may occur within minutes of carboplatin administration. Epinephrine, corticosteroids, and antihistamines have been employed to alleviate symptoms

DESCRIPTION:

Carboplatin Injection is supplied as a sterile, pyrogen-free solution available in 10 mg per mL multiple dose vials containing 50 mg. 150 mg, 450 mg or 600 mg of carboplatin for administration by intravenous infusion. Each mL contains: carboplatin 10 mg, and water for injection to volume.

Carboplatin is a platinum coordination compound. The chemical name for carboplatin is platinum, diammine [1,1-cyclobutanedicarboxylato(2-)-0,0']-, (SP-4-2), and has the following structural formula:



451023D /Revised: May 2021

fully.

entire leaflet carteren

this e it for

ad ep

platin

BO

2 ectio only

D

Information

Patient

CARBOplatin Rx only

FRESENIUS

KABI

 $C_6H_{12}N_2O_4Pt$ Carboplatin is a crystalline powder. It is soluble in water at a rate of approximately 14 mg/mL, and the pH of a 1% solution is 5 to 7 It is virtually insoluble in ethanol, acetone, and dimethylacetamide. CLINICAL PHARMACOLOGY:

Carboplatin, like cisplatin, produces predominantly interstrand DNA cross-links rather than

DNA-protein cross-links. This effect is apparently cell-cycle nonspecific. The aquation of carboplatin, which is thought to produce the active species, occurs at a slower rate than in the case of cisplatin. Despite this difference, it appears that both carboplatin and cisplatin induce equal numbers of drug-DNA cross-links, causing equivalent lesions and biological effects. The differences in potencies for carboplatin and cisplatin appear to be directly related to the difference in aquation

In patients with creatinine clearances of about 60 mL/min, or greater, plasma levels of intact carboplatin decay in a biphasic manner after a 30-minute intravenous infusion of 300 to 500 mg/m² of carboplatin. The initial plasma half-life (alpha) was found to

o you learn more ction. It cannot, ossible warnings to Carboplatin ot list all of the arboplatin Injec-always be your formation about and your treat-ur doctor about

ot to

ЪС

n will help atin Inject all the pos

δO

.⊆ t∃

it does not l isks of Carbo tor should alv detailed inforr condition an to ask your o you may hav

refits and rist refits and rist renoice for d ur medical c rnt. Be sure t questions y

c det is k

/ever, cov side effec ction, an lefits and . Your do choice fo

be 1.1 to 2 hours (n=6), and the post distribution plasma half-life (beta) was found to be 2.6 to 5.9 hours (n=6). The total body clearance, apparent volume of distribution and mean residence time for carboplatin are 4.4 L/hour. 16 L and 3.5 hours. respectively The C_{may} values and areas under the plasma concentration versus time curves from 0 to infinity (AUC inf) increase linearly with dose, although the increase was slightly more than dose proportional. Carboplatin, therefore, exhibits linear pharmacokinetics over the dosing range studied (300 to 500 mg/m²).

Carboplatin is not bound to plasma proteins. No significant quantities of proteinfree, ultrafilterable platinum-containing species other than carboplatin are present in plasma. However, platinum from carboplatin becomes irreversibly bound to plasma proteins and is slowly eliminated with a minimum half-life of 5 days.

The major route of elimination of carboplatin is renal excretion. Patients with creatinine clearances of approximately 60 mL/min or greater excrete 65% of the dose in the urine within 12 hours and 71% of the dose within 24 hours. All of the platinum in the 24-hour urine is present as carboplatin. Only 3% to 5% of the administered platinum is excreted in the urine between 24 and 96 hours. There are insufficient data to determine whether biliary excretion occurs.

In patients with creatinine clearances below 60 mL/min, the total body and renal clearances of carboplatin decrease as the creatinine clearance decreases. Carboplatin dosages should therefore be reduced in these patients (see DOSAGE AND ADMINISTRATION).

The primary determinant of carboplatin clearance is glomerular filtration rate (GFR) and this parameter of renal function is often decreased in elderly patients. Dosing formulas incorporating estimates of GFR (see DOSAGE AND ADMINISTRATION) to provide predictable carboplatin plasma AÚCs should be used in elderly patients to minimize the risk of toxicity.

CLINICAL STUDIES:

Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer

In two prospectively randomized, controlled studies conducted by the National Cancer Institute of Canada, Clinical Trials Group (NCIC) and the Southwest Oncology Group (SWOG), 789 chemotherapy naive patients with advanced ovarian cancer were treated with carboplatin or cisplatin, both in combination with cyclophosphamide every 28 days for 6 courses before surgical reevaluation. The following results were obtained from both studies:

Comparative Efficacy Overview of Pivotal Trials SMUG

Number of patients randomized	447	342
Median age (years)	60	62
Dose of cisplatin	75 mg/m ²	100 mg/m ²
Dose of carboplatin	300 mg/m ²	300 mg/m ²
Dose of cyclophosphamide	600 mg/m ²	600 mg/m ²
Residual tumor <2 cm	39% (174/447)	14% (49/342)
(number of patients)	. ,	. ,

Comparative Efficacy (Continued) ahle Diseasé Patient

	NCIC	SWOG
Carboplatin (number of patients)	60% (48/80)	58% (48/83)
Čisplatin .	58% (49/85)	43% (33/76)
(number of patients) 95% C.I. of difference (Carboplatin–Cisplatin)	(-13.9%, 18.6%)	(-2.3%, 31.1%)
Pathologic	Complete Respons	se* SWOG

Carboplatir

11% (24/224) 10% (17/171) (number of patients) 10% (17/171) 15% (33/223) (number of patients) 95% C L of difference (-10.7%, 2.5%) (-6.9%, 6.9%) (Carboplatin–Cisplatin) *114 Carboplatin and 109 Cisplatin patients did not undergo 90 Carboplatin and 100 Singlatin patients of 100 undergo second look surgery in NCIC study.
 90 Carboplatin and 106 Cisplatin patients did not undergo second look surgery in SWOG study. Progression-Free Survival (PFS) NCIC ŚWOG Carboplati 59 weeks 49 weeks Cisplatin 2-year PFS* 61 weeks 47 weeks 31% Cisplatin 95% C.I. of difference (-9.0, 9.4) (-9.3, 8.7)

(Carboplatin-Cisplatin) 3-year PFS* Cisolatin 95% C.I. of difference (-11.5, 4.5) (-14.1, 0.3) (Carboplatin-Cisplatin) Hazard Ratio* 95% C.I. 1 10 (0.89, 1.35) (0.81, 1.29) (Carboplatin-Cisplatin) *Kaplan-Meier Estimates Unrelated deaths occurring in the absence of progression were counted as events (progression) in this analysis. *Analysis adjusted for factors found to be of prognostic significance were consistent with unadjusted analysis.

	Survival NCIC	SWOG
Median		
Carboplatin	110 weeks	86 weeks
Cisplatin	99 weeks	79 weeks
2-vear Survival*		
Carboplatin	51.9%	40.2%
Cisplatin	48.4%	39%
95% C.I. of difference	(-6.2, 13.2)	(-9.8, 12.2)
(Carboplatin-Cisplatin)	· · /	(, ,
3-year Survival*		
Carboplatin	34.6%	18.3%
Cisplatin	33.1%	24.9%
95% C.I. of difference	(-7.7, 10.7)	(-15.9, 2.7)
(Carboplatin-Cisplatin)	· · /	(, ,
Hazard Ratio**		
95% C.I.	0.98	1.01
(Carboplatin-Cisplatin)	(0.78, 1.23)	(0.78, 1.30)
*Kaplan Major Estimate		

*Kaplan-Meier Estimates *Analysis adjusted for factors found to be of prognostic significance were consistent with unadjusted analysis.

Comparative Toxicity

The pattern of toxicity exerted by the carboplatin-containing regimen was significantly different from that of the cisplatin-containing combinations. Differences between the two studies may be explained by different cispl-atin dosages and by different supportive care. The carboplatin-containing regimen induced significantly more thrombocytopenia and, in one study, significantly more leukopenia and more need for transfusional support. The cisplatin-containing regimen produced significantly more anemia in one study. However, no significant differences occurred in incidences of infections and hemorrhagic episodes.

Non-hematologic toxicities (emesis, neurotoxicity, ototoxicity, renal toxicity, hypomagnesemia, and alopecia) were significantly more frequent in the cisplatin-containing

Bone Marrow Thrombocytopenia <100,000/mm³ 70 29 <1 <50,000/mm³ 41 6 <1 Neutropenia < <1 < 2,000 cells/mm³ 97 96 <1,000 cells/mm³ 81 79 Leukopenia < <4,000 cells/mm³ 98 97	alues** 0.001 0.001 n.s. n.s. 0.001
Arm Arm Percent* Percent* P-V Bone Marrow Thrombocytopenia <100,000/mm³ 70 29 <i< td=""> <50,000/mm³ 70 29 <i< td=""> <50,000/mm³ 41 6 <i< td=""> Neutropenia <i< td=""> <2,000 cells/mm³ 97 96 <i,000 cells="" mm³<="" td=""> <4,000 cells/mm³ 81 79 Leukopenia <4,000 cells/mm³ 98 97 <z,000 cells="" mm³<="" td=""> 68 52 C Anemia</z,000></i,000></i<></i<></i<></i<>	0.001 0.001 n.s. n.s. n.s.
Thrombocytopenia <100,000/mm³ 70 29 <1 <50,000/mm³ 41 6 <1 Neutropenia 2 <1 6 <1 < 2,000 cells/mm³ 97 96 <1,000 cells/mm³ 81 79 Leukopenia <1 68 52 C Anemia	0.001 n.s. n.s. n.s.
Thrombocytopenia <100,000/mm³	0.001 n.s. n.s. n.s.
<pre><100,000/mm³ 70 29 <1 <pre><1 <pre><100,000/mm³ 41 6 <1 Neutropenia <pre><2,000 cells/mm³ 97 96 <1,000 cells/mm³ 81 79 Leukopenia <4,000 cells/mm³ 98 97 <2,000 cells/mm³ 68 52 0 Anemia</pre></pre></pre></pre>	0.001 n.s. n.s. n.s.
Neutropenia 97 96 < 2,000 cells/mm³	n.s. n.s. n.s.
< 2,000 cells/mm ³ 97 96 <1,000 cells/mm ³ 81 79 Leukopenia <4,000 cells/mm ³ 98 97 <2,000 cells/mm ³ 68 52 (0 Anemia	n.s. n.s.
<pre><1,000 cells/mm³ 81 79 Leukopenia <4,000 cells/mm³ 98 97 <2,000 cells/mm³ 68 52 0 Anemia</pre>	n.s. n.s.
Leukopenia <4,000 cells/mm ³ 98 97 <2,000 cells/mm ³ 68 52 (Anemia	n.s.
<4,000 cells/mm ³ 98 97 <2,000 cells/mm ³ 68 52 0 Anemia	
<2,000 cells/mm ³ 68 52 0 Anemia	
Anemia	
	n.s.
<8 g/dL 18 12	n.s.
	n.s.
	n.s.
Transfusions 42 31 0	0.018
Gastrointestinal	
· · · · · · · · · · · · · · · ·	0.010
	0.001
Other GI side effects 50 62 0	0.013
Neurologic	
	0.001
	0.001
	n.s.
effects Central neurotoxicity 28 40 0	0.009
	1.009
Renal	
	0.006
elevations Blood urea elevations 17 31 <(0.001
	0.001
Hepatic	
	n.s. n.s.
Alkaline phosphatase – –	-
elevations	
Electrolytes loss	
	0.005
	n.s.
	n.s.
	0.001
Other side effects	
	n.s.
	n.s.).017
	n.s.

*Values are in percent of evaluable patients.

**n.s. = not significant, p > 0.05.
 *May have been affected by cyclophosphamide dosage delivered.

-forming ssion) or ē oth me pr baby

őδ

give ent j How is Carboplatin Injection us how is Carboplatin Injection us only a professional experienced in se of cancer drugs should give is medication. Carboplatin Injectic jiven by dripping the medicine slc ind directly into a vein (intraven ritusion) for 15 minutes or lon (our doctor will determine the dos carboplatin Injection for you based our weight, height, and kidney fu on. Carboplatin Injection may be g lone or with other drugs. Treatmel sually repeated every four weeks f umber of cycles. Before and after Carboplatin Injec reatment, your doctor may give nedication to lessen the nausea omiting associated with this car be c be c atme

stion you and cer latin Injec nay give nausea n this car

What is cancer? What is cancer? Under normal conditions, the cells in your body divide and grow in an orderly, controlled fashion. Cell division and growth are necessary for the human body to perform its functions and to repair itself. Cancer cells are different from normal cells because they are not able to control their own growth. The reasons for this abnormal growth are not yet fully understood. A tumor is a mass of unhealthy cells that are dividing and growing fast and in an uncontrolled way. When a tumor invades surrounding healthy body tissue it is known as a malignant tumor. A malignant tumor can spread (metasta-size) from its original location to other parts of the body.

althy cells j fast and n a tumor ody tissue tumor. A (metasta-n to other

문 Φ e e e ÷ ≥ŏ rboplatin I Injection i reat cancentering nuttiplying ວວ is Carb pplatin I ed to tre s by int pidly mu er cells. e ti te te acta

arboplatin Ű take not ould o sh ctior 2

Injection latin Ido N ਰ ਜ਼ ਸ਼ੁ ded i to car with

<u>.</u>0

- ٥ platin ned ned nen gic con atment recomm re aller latinum ave a ystem (b
- lood-1 depres ≤ ه Ð
- to becon eding a l (bone ant blee jnant, i are br or . بتو

ADVERSE EX WITH OVARIA	AN CANCER	SWOG ST	
(Carboplatin Arm	Cisplatin Arm	
			P-Values**
arrow			
ocytopenia			
0,000/mm ³	59	35	< 0.001
),000/mm³ enia	22	11	0.006
000 cells/mm ³	95	97	n.s.
100 cells/mm ³ enia	84	78	n.s.
000 cells/mm ³	97	97	n.s.
000 cells/mm ³	76	67	n.s.
g/dL	88	87	n.s.
/dL	8	24	< 0.001
ns	18	21	n.s.
9	6	4	n.s.
sions	25	33	n.s.
ntestinal			
and vomiting		96	n.s.
g	82	91	0.007
l side effects	40	48	n.s.
gic			
ral neuropathie		28	0.001
city	12	30	< 0.001
ensory side ts	4	6	n.s.
neurotoxicity	23	29	n.s.
creatinine tions	7	38	<0.001
rea elevations	-	-	-
, elevations	5	3	n.s.
levations	23	16	n.s.
phosphatase tions		20	n.s.
ytes loss			
	-	-	-
um	_	-	-
ium	58	77	< 0.001
ide effects			
	54	52	n.s.
a	43	46	n.s.
ascular	23	30	n.s.
tory	12	11	n.s.
	10	11	n.s.
rinary	11	13	n.s.
a+	43	57	0.009
tis	6	11	n.s.

Bone Marrow

tropenia

Leukopenia

<11 g/d

<8 g/dL

Transfusions

Gastrointestinal

Nausea and vomiting

Other GI side effects

Other sensory side

Serum creatinine

Bilirubin elevations

SGOT elevations

elevations

Electrolytes loss

Other side effects

elevations

Hepatic

Sodium

Potassium

Magnesium

Asthenia

Cardiovascular

Genitourinary

**n.s. = not significant. p > 0.05.

Alopecia+

Mucositis

71+ weeks

Respiratory

Alleraic

Calcium

Central neurotoxicity

Blood urea elevations

Alkaline phosphatase

Peripheral neuropathies 13

Infections

Bleedina

Vomitina

Neurologic

Ototoxicity

effects

Thrombocytopenia

<100,000/mm

< 50,000/mm

<2.000 cells/mm³

<1.000 cells/mm³

<4,000 cells/mm³

<2,000 cells/mm³

*Values are in percent of evaluable patients.

*May have been affected by cyclophosphamide dosage delivered.

Use as a Single Agent for Secondary Treatment of Advanced Ovarian Cancer In two prospective, randomized controlled studies in patients with advanced ovarian cancer previously treated with chemotherapy carboplatin achieved 6 clinical complete responses in 47 patients. The duration of these responses ranged from 45 to

INDICATIONS: Initial Treatment of Advanced Ovarian Carcinoma

Carboplatin Injection is indicated for the initial treatment of advanced ovarian carcinoma in established combination with other approved chemotherapeutic agents. One established combination regimen consists of carboplatin and cyclophosphamide. Two randomized controlled studies conducted by the NCIC and SWOG with carboplatin versus cisplatin both in combination with cyclophosphamide have demonstrated equivalent overall survival between the two groups (see CLINICAL STUDIES)

There is limited statistical power to demon strate equivalence in overall pathologic complete response rates and long-term survival (≥3 years) because of the small number of patients with these outcomes the small number of patients with residual tumor <2 cm after initial surgery also limits the statistical power to demonstrate equivalence in this subgroup.

Secondary Treatment of Advanced Ovarian Carcinoma

Carboplatin Injection is indicated for the palliative treatment of patients with ovarian carcinoma recurrent after prior chemotherapy, including patients who have been previously treated with cisplatin.

Within the group of patients previously treated with cisplatin, those who have devel oped progressive disease while receiving cisplatin therapy may have a decreased response rate

CONTRAINDICATIONS:

Carboplatin Injection is contraindicated in patients with a history of severe allergic reactions to cisplatin or other platinum-containing compounds

Carboplatin Injection should not be employed in patients with severe bone marrow depression or significant bleeding.

WARNINGS:

Bone marrow suppression (leukopenia neutropenia, and thrombocytopenia) is dose-dependent and is also the dose-limiting toxicity. Peripheral blood counts should be frequently monitored during carboplatin treatment and, when appropriate, until recovery is achieved. Median nadir occurs at day 21 in patients receiving single-agent carboplatin. In general, single intermittent courses of carboplatin should not be repeated until leukocyte, neutrophil, and platelet counts have recovered

Since anemia is cumulative, transfusions may be needed during treatment with carboplatin, particularly in patients receiving prolonged therapy.

Bone marrow suppression is increased in patients who have received prior therapy especially regimens including cisplatin Marrow suppression is also increased in patients with impaired kidney function. Initial carboplatin dosages in these patients should be appropriately reduced (see DOSAGE AND ADMINISTRATION) and blood counts should be carefully monitored between

courses. The use of carboplatin in combination with other bone marrow suppressing therapies must be carefully managed with respect to dosage and timing in order to minimize additive effects.

Carboplatin has limited nephrotoxic potential, but concomitant treatment with aminoglycosides has resulted in increased renal and/or audiologic toxicity, and caution must be exercised when a patient receives both drugs. Clinically significant hearing loss has been reported to occur in pediatric patients when carboplatin was administered at higher than recommended doses in combination with other ototoxic agents.

Carboplatin can induce emesis, which can be more severe in patients previously receiving emetogenic therapy. The incidence and intensity of emesis have been reduced by using premedication with antiemetics. Although no conclusive efficacy data exist with the following schedules of carboplatin, lengthening the duration of single intravenous administration to 24 hours or dividing the total dose over 5 consecutive daily pulse doses has resulted in reduced emesis.

Although peripheral neurotoxicity is infrequent, its incidence is increased in patients older than 65 years and in patients previously treated with cisplatin. Pre-existing cisplatininduced neurotoxicity does not worsen in about 70% of the patients receiving carboplatin as secondary treatment.

Loss of vision, which can be complete for light and colors, has been reported after the use of carboplatin with doses higher than those recommended in the package insert. Vision appears to recover totally or to a significant extent within weeks of stopping these high doses.

As in the case of other platinum-coordination compounds, allergic reactions to carboplatin have been reported. These may occur within minutes of administration and should be managed with appropriate supportive therapy. There is increased risk of allergic reactions including anaphylaxis in patients previously exposed to platinum therapy (see CONTRAINDICATIONS and ADVERSE REACTIONS: Allergic Reactions).

High dosages of carboplatin (more than 4 times the recommended dose) have resulted in severe abnormalities of liver function tests.

Carboplatin Injection may cause fetal harm when administered to a pregnant woman. Carboplatin has been shown to be embryotoxic and teratogenic in rats. There are no adequate and well-controlled studies in pregnant women. If this drug is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant.

PRECAUTIONS: Genera

Needles or intravenous administration sets containing aluminum parts that may come in contact with Carboplatin Injection should not be used for the preparation or administration of the drug. Aluminum can react with carboplatin causing precipitate formation and loss of potency

Drug Interactions

The renal effects of nephrotoxic compounds may be potentiated by carboplatin.

Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of carboplatin has not been studied, but compounds with similar mechanisms of action and mutagenicity profiles have been reported to be carcinogenic. Carboplatin has been shown to be mutagenic both in vitro and in vivo. It has also been shown to be embryotoxic and teratogenic in rats receiving the drug during organogenesis. Secondary malignan-cies have been reported in association with multi-drug therapy.

Nursing Mothers

It is not known whether carboplatin is excreted in human milk. Because there is a possibility of toxicity in nursing infants secondary to carboplatin treatment of the mother, it is recommended that breast-feeding be discontinued if the mother is treated with Carboplatin Injection.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established (see WARNINGS; "audiologic toxicity")

Geriatric Use

Of the 789 patients in initial treatment combination therapy studies (NCIC and SWOG), 395 patients were treated with carboplatin in combination with cyclophosphamide Of these, 141 were over 65 years of age and 22 were 75 years or older. In these trials, age was not a prognostic factor for survival. In terms of safety, elderly patients treated with carboplatin were more likely to develop severe thrombocytopenia than younger patients. In a combined database of 1,942 patients (414 were ≥65 years of age) that received single-agent carboplatin for different tumor types, a similar incidence of adverse events was seen in patients 65 years and older and in patients less than 65. Other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Because renal function is often decreased in the elderly, renal function should be considered in the selection of carboplatin dosage (see DOSAGE AND ADMINISTRATION).

ADVERSE REACTIONS:

To report SUSPECTED ADVERSE REAC-TIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

For a comparison of toxicities when carboplatin or cisplatin was given in combination with cyclophosphamide, see CLINICAL STUDIES, Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer. Comparative Toxicity

What should you tell your doctor before starting treatment with Carboplatin Injection?
Discuss the benefits and risks of Carboplatin Injection with your doctor before beginning treatment.
Be sure to inform your doctor before beginning treatment.
If you are allergic to carboplatin or other platinum-containing products;
If you are or intend to become pregnant, since carboplatin may harm the developing fetus. It is important to use being treated with Carboplatin life you are breast-feeding, since nursing infants may be exposed to carboplatin in this way;

- icines d non drugs actior and the ption affect ou are taking othe uding all prescript scription (over-the-co e carboplatin may al ther medicines; ou have any other r s, especially chicken ant exposure to adu in chicken pox), shir ofens, infection, or k othe /ou sscri othe c effection of concentration of concentrat
 - pro i<u>c</u> i cal ゔ゠ゔ medic pox (i ults or ngles, n po) lults kidn e to ox), tion, you ms, €

king tal

þe might l feedin /. Carb **d l avoid while tal injection?** Inant or think you migh if you are breast feed r know right wawy. Ca on may harm your de or not childeding bat man of childearing se birth control to a aint while you are ta avoid contact with ac who have infections, or right away if you s fion such as cough, ff Also, while you are b arboplatin Injection or arboplatin Injection or arboplatin Check with re to a re to a s, while t in? ithink you are bre? vright t-' What should I Carboplatin In If you are pregnan pregnant, or if y platin Injection i you should use you should use you should use you should use tell your doctor i signs of infectior signs of infectior signs of infectior out out doctor i signs of infectior out out the street you stop treatme

...e t Jon or Jon or Jon or Jon or Vithey c they c oplatin Inj nt, first ch Jetting ar Is). Avoic 2 e e wh. ng le 9 id ation us ac <u>ح د ج</u> s (v poli

Cal

e the possible side effects oplatin Injection? latin Injection may cause d effects, particularly because atin Injection interferes with the f normal cells as well as cancer or example, the occurrence of cancer (secondary malignancy) odated

ADVERSE EXPERIENCES IN PATIENTS WITH OVARIAN CANCER		
	First Line Combination Therapy* Percent	Second Line Single Agent Therapy** Percent
Bone Marrow		
Thrombocytopenia		
<100,000/mm ³	66	62
< 50,000/mm ³	33	35
Neutropenia		
<2,000 cells/mm ³	96	67
<1,000 cells/mm ³	82	21
Leukopenia		
<4,000 cells/mm ³	97	85
<2,000 cells/mm ³	71	26
Anemia		
<11 g/dL	90	90
<8 g/dL	14	21
Infections	16	5
Bleeding	8	5
Transfusions	35	44
Gastrointestinal		
Nausea and vomiting	93	92
Vomiting	83	81
Other GI side effects	46	21
Veurologic		
Peripheral neuropathies	15	6
Dtotoxicity	12	ĭ
Other sensory side effects	5	1
Central neurotoxicity	26	5
Renal		-
Serum creatinine elevations	6	10
Blood urea elevations	17	22
	17	22
Hepatic	-	-
Bilirubin elevations	5	5
SGOT elevations	20	19
Alkaline phosphatase elevation	ns 29	37
Electrolytes loss		
Sodium	10	47
Potassium	16	28
Calcium	16	31
Magnesium	61	43
Other side effects		
Pain	44	23
Asthenia	41	11
Cardiovascular	19	6
Respiratory	10	6
Allergic	11	ž
Genitourinary	10	2
Alopecia	49	2
Mucositis	8	1
	~	· ·

*Use with Cyclophosphamide for Initial Treatment of Ovarian Cancer

Data are based on the experience of 393 patients with ovarian cancer (regardless of baseline status) who received initial combination therapy with carboplatin and cyclo phosphamide in two randomized controlled studies conducted by SWOG and NCIC (see CLINICAL STUDIES

Combination with cyclophosphamide as well as duration of treatment may be responsible for the differences that can be noted in the adverse experience table.

**Single Agent Use for the Secondary Treatment of Ovarian Cancer

Data are based on the experience of



553 patients with previously treated ovarian carcinoma (regardless of baseline status) who received single-agent carboplatin.

In the narrative section that follows, the incidences of adverse events are based on data from 1.893 patients with various types of tumors who received carboplatin as singleagent therapy.

Hematologic Toxicity

Bone marrow suppression is the dose-limiting toxicity of carboplatin. Thrombocytopenia with platelet counts below 50.000/mm³ occurs in 25% of the patients (35% of pretreated ovarian cancer patients); neutropenia with granulocyte counts below 1,000/mm³ occurs in 16% of the patients (21% of pretreated ovarian cancer patients): leukopenia with WBC counts below 2 000/mm³ occurs in 15% of the patients (26% of pretreated ovarian cancer patients). The nadir usually occurs about day 21 in patients receiving single agent therapy. By day 28, 90% of patients have platelet counts above 100 000/mm³ 74% have neutrophil counts above 2 000/mm³ 67% have leukocyte counts above 4 000/mm³

Marrow suppression is usually more severe in patients with impaired kidney function. Patients with poor performance status have also experienced a higher incidence of severe leukopenia and thrombocytopenia.

The hematologic effects, although usually reversible, have resulted in infectious or hemorrhagic complications in 5% of the patients treated with carboplatin, with drugrelated death occurring in less than 1% of the patients. Fever has also been reported in patients with neutropenia.

Anemia with hemoglobin less than 11 g/dL has been observed in 71% of the patients who started therapy with a baseline above that value. The incidence of anemia increases with increasing exposure to carboplatin. Transfusions have been administered to 26% of the patients treated with carboplatin (44% of previously treated ovarian cancer patients).

Bone marrow depression may be more severe when carboplatin is combined with other bone marrow suppressing drugs or with radiotherapy.

Gastrointestinal Toxicity

Vomiting occurs in 65% of the patients (81% of previously treated ovarian cancer patients) and in about one-third of these patients it is severe. Carboplatin, as a single agent or in combination, is significantly less emetogenic than cisplatin; however, patients previously treated with emetogenic agents, especially cisplatin, appear to be more prone to vomiting. Nausea alone occurs in an additional 10% to 15% of patients. Both nausea and vomiting usually cease within 24 hours of treatment and are often responsive to antiemetic measures. Although no conclusive efficacy data exist with the following schedules, prolonged administration of carboplatin, either by continuous 24-hour infusion or by daily pulse doses given for 5 consecutive days, was associated with less severe vomiting than the single dose intermittent schedule. Emesis was increased when carboplatin was used in combination with other emetogenic compounds. Other gastrointestinal effects observed frequently were pain, in 17% of the patients; diarrhea in 6%: and constipation, also in 6%.

Neurologic Toxicity

Peripheral neuropathies have been observed in 4% of the patients receiving carboplatin (6% of pretreated ovarian cancer patients) with mild paresthesias occurring most frequently. Carboplatin therapy produces significantly fewer and less severe neurologic side effects than does therapy with cisplatin However, patients older than 65 years and/or previously treated with cisplatin appear to have an increased risk (10%) for peripheral neuropathies. In 70% of the patients with pre-existing cisplatin-induced peripheral neurotoxicity, there was no worsening o symptoms during therapy with carboplatin. Clinical ototoxicity and other sensory abnor malities such as visual disturbances and change in taste have been reported in only 1% of the patients. Central nervous system symptoms have been reported in 5% of the patients and appear to be most often related to the use of antiemetics.

Although the overall incidence of peripheral neurologic side effects induced by carboplatin is low, prolonged treatment, particularly in cisplatin pretreated patients, may result in cumulative neurotoxicity.

Nephrotoxicity

Development of abnormal renal function test results is uncommon despite the fact that carboplatin, unlike cisplatin, has usually been administered without high-volume fluid hydration and/or forced diuresis. The incidences of abnormal renal function tests reported are 6% for serum creatinine and 14% for blood urea nitrogen (10% and 22% respectively in pretreated ovarian cancer patients). Most of these reported abnormalities have been mild and about one-half of them were reversible Creatinine clearance has proven to be the most sensitive measure of kidney function in patients receiving carboplatin, and it appears to be the most useful test for correlating drug clearance and bone marrow suppression Twenty-seven percent of the patients who had a baseline value of 60 mL/min or more demonstrated a reduction below this value during carboplatin therapy.

Hepatic Toxicity

The incidences of abnormal liver function tests in patients with normal baseline values were reported as follows: total bilirubin, 5%; SGOT, 15%; and alkaline phosphatase, 24%; (5%, 19%, and 37%, respectively, in pretreated ovarian cancer patients). These abnormalities have generally been mild and reversible in about one-half of the cases, although the role of metastatic tumor in the liver may complicate the assessment in many patients. In a limited series of patients receiving very high dosages of carboplatin and autologous bone marrow transplantation, severe abnormalities of liver function tests were reported.

Electrolyte Changes

The incidences of abnormally decreased serum electrolyte values reported were as follows: sodium, 29%; potassium, 20% calcium, 22%; and magnesium, 29%; (47% 28% 31% and 43% respectively in pretreated ovarian cancer patients). Electrolyte supplementation was not routinely administered concomitantly with carboplatin, and these electrolyte abnormalities were rarely associated with symptoms.

Allergic Reactions

Hypersensitivity to carboplatin has been reported in 2% of the patients. These allergic reactions have been similar in nature and severity to those reported with other platinumcontaining compounds, ie. rash, urticaria. erythema, pruritus, and rarely bronchospasm and hypotension. Anaphylactic reactions have been reported as part of postmarketing surveillance (see WARNINGS). These reac tions have been successfully managed with standard epinephrine corticosteroid and antihistamine therapy

Injection Site Reactions

Injection site reactions, including redness. swelling, and pain, have been reported during postmarketing surveillance. Necrosis associated with extravasation has also been reported

Other Events

Pain and asthenia were the most frequently reported miscellaneous adverse effects: thei relationship to the tumor and to anemia was likely Alopecia was reported (3%) Cardio vascular, respiratory, genitourinary, and mucosal side effects have occurred in 6% or less of the patients. Cardiovascular events (cardiac failure, embolism, cerebrovascular accidents) were fatal in less than 1% of the patients and did not appear to be related to chemotherapy. Cancer-associated hemolytic uremic syndrome has been reported rarely. Malaise, anorexia, hypertension, dehydra

tion, and stomatitis have been reported as part of postmarketing surveillance.

OVERDOSAGE:

There is no known antidote for Carboplatin Injection overdosage. The anticipated complications of overdosage would be secondary to bone marrow suppression and/or hepatic toxicity.

DOSAGE AND ADMINISTRATION:

NOTE: Aluminum reacts with carboplatin causing precipitate formation and loss of potency, therefore, needles or intravenous sets containing aluminum parts that may come in contact with the drug must not be used for the preparation or administration of Carboplatin Injection.

Single-Agent Therapy

Carboplatin Injection, as a single agent, has been shown to be effective in patients with recurrent ovarian carcinoma at a dosage of 360 mg/m² IV on day 1 every 4 weeks (alternatively see Formula Dosing). In general, however, single intermittent courses of

Carboplatin Injection should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100.000.

Combination Therapy with Cvclophosphamide

In the chemotherapy of advanced ovarian cancer, an effective combination for previously untreated patients consists of:

Carboplatin Injection-300 mg/m² IV on day 1 every 4 weeks for 6 cycles (alternatively see Formula Dosing).

Cyclophosphamide-600 mg/m2 IV on day 1 every 4 weeks for 6 cycles. For directions regarding the use and administration of cyclophosphamide please refer to its package insert (see CLINICAL STUDIES). Intermittent courses of Carboplatin Injec-

tion in combination with cyclophosphamide should not be repeated until the neutrophil count is at least 2,000 and the platelet count is at least 100 000

Dose Adjustment Recommendations

Pretreatment platelet count and performance status are important prognostic factors for severity of myelosuppression in previously treated patients

The suggested dose adjustments for single agent or combination therapy shown in the table below are modified from controlled trials in previously treated and untreated patients with ovarian carcinoma. Blood counts were done weekly, and the recommendations are based on the lowest post-treatment platelet or neutrophil value.

Platelets	Neutrophils	Adjusted Dose* (From Prior Course)
>100,000	>2000	125%
50 to 100,000	500 to 2000	No Adjustment
<50,000	<500	75%

*Percentages apply to Carboplatin Injection as a single

agent or to both carboplatin and cyclophosphamide in nbination. In the controlled studies, dosages were also adjusted at a lower level (50 to 60%) for severe myelosuppression. Escalations above 125% were not recommended for these studies.

Carboplatin Injection is usually administered by an infusion lasting 15 minutes or longer. No pre- or post-treatment hydration or forced diuresis is required.

Patients with Impaired Kidney Function Patients with creatinine clearance values below 60 mL/min are at increased risk of severe bone marrow suppression. In renallyimpaired patients who received single-agent carboplatin therapy, the incidence of severe leukopenia, neutropenia, or thrombocyto penia has been about 25% when the dosage modifications in the table below have been used

	Baseline	
Creatinine Clearance		Recommended Dose on Day 1
41 to 59 mL/min 16 to 40 mL/min		250 mg/m ² 200 mg/m ²

The data available for patients with severely impaired kidney function (creatinine clearance below 15 mL/min) are too limited to permit a recommendation for treatment. These dosing recommendations apply to

> Adm Adm Patient Inform roved by the ninistration. ⊂ब् νģ ו Leaflet Food נ and Drug

can repl doctor. tion upon 4. You may 1 with your summary with your

the initial course of treatment. Subsequent dosages should be adjusted according to the patient's tolerance based on the degree of bone marrow suppression.

Formula Dosing

of developing toxicity.

risk of toxicity.

SOLUTIONS:

Chloride Injection, USP

Geriatric Dosing

Another approach for determining the initial dose of Carboplatin Injection is the use of mathematical formulae, which are based on a patient's pre-existing renal function or renal function and desired platelet nadir. Renal excretion is the major route of elimination for carboplatin (see CLINICAL PHARMA-COLOGY). The use of dosing formulae, as compared to empirical dose calculation based on body surface area, allows compensation for patient variations in pretreatment renal function that might otherwise result in either underdosing (in patients with above average renal function) or overdosing (in patients with impaired renal function).

A simple formula for calculating dosage based upon a patient's glomerular filtration rate (GFR in mL/min) and Carboplatin Injection target area under the concentration versus time curve (AUC in mg/mL•min), has been proposed by Calvert. In these studies, GFR was measured by ⁵¹Cr-EDTA clearance.

CALVERT FORMULA FOR CARBOPLATIN DOSING Total Dose (mg) = (target AUC) x (GFB + 25) Note: With the Calvert formula, the total dose of carboplatin is calculated in mg, not mg/m².

The target AUC of 4 mg/mL·min to 6 mg/mL•min using single-agent carboplatin appears to provide the most appropriate dose range in previously treated patients. This study also showed a trend between the AUC of single-agent carboplatin administered to previously treated patients and the likelihood

% Actual	Toxicity in Previously Gr 3 or Gr 4	Treated Patients Gr 3 or Gr 4
AUC (mg/mL•min)	Thrombocytopenia	Leukopenia
4 to 5	16%	13%
6 to 7	33%	34%

Because renal function is often decreased in elderly patients, formula dosing of carboplatin based on estimates of GFR should be used in elderly patients to provide predictable plasma carboplatin AUCs and thereby minimize the

PREPARATION OF INTRAVENOUS

Carboplatin Injection is a premixed aqueous solution of 10 mg/mL carboplatin.

Carboplatin Injection can be further diluted to concentrations as low as 0.5 mg/mL with 5% Dextrose in Water (D₅W) or 0.9% Sodium

When prepared as directed, Carboplatin Injection solutions are stable for 8 hours at room temperature (25°C). Since no antibacterial preservative is contained in the formulation. it is recommended that Carboplatin Injection solutions be discarded 8 hours after dilution

HOW SUPPLIED: Product NDC No. No

107260 63323-172-60 CARBOplatin Injection, 600 mg per 60 mL (10 mg per mL), in a 60 mL multiple dose vial packaged individually

The container closure is not made with natural rubber latex.

Storage

Unopened vials of Carboplatin Injection are stable to the date indicated on the package when stored at 25°C (77°F); [excursions permitted from 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature] Protect from light.

Carboplatin Injection multidose vials maintain microbial, chemical, and physical stability for up to 14 days at 25°C following multiple needle entries.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Solutions for infusion should be discarded 8 hours after preparation.

Handling and Disposal

Caution should be exercised in handling and preparing carboplatin injection. Severa guidelines on this subject have been nublished 1-4

To minimize the risk of dermal exposure always wear impervious gloves when handling vials containing carboplatin injection. If carboplatin injection contacts the skin, immediately wash the skin thoroughly with soap and water. If carboplatin inje tion contacts mucous membranes, the membranes should be flushed immediately and thoroughly with water. More information is available in the references listed below.

REFERENCES:

- 1. NIOSH Alert: Preventing occupational exposures to antineoplastic and other hazardous drugs in healthcare settings. 2004. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2004-165.
- 2. OSHA Technical Manual, TED 1-0.15A Section VI: Chapter 2. Controlling occupational exposure to hazardous drugs. OSHA, 1999. http://www.osha.gov/dts/osta/otm/otm_vi/ otm vi 2 html
- 3. American Society of Health-System Pharmacists. ASHP guidelines on handling hazardous drugs. Am J Health-Syst Pharm. 2006;63: 1172-1193.
- 4. Polovich M. White JM. Kelleher LO. eds 2005. Chemotherapy and biotherapy guidelines and recommendations to practice. 2nd ed. Pittsburgh, PA: Oncology Nursing Society.



www.fresenius-kabi.com/us

451023D Revised: May 2021

This medicine was prescribed for your particular condition. It must be given under close medical supervision by a doctor trained in the use of drugs for the treatment of cancer. This summary does not include every-thing there is to know about Carboplatin Injection. Medicines are sometimes prescribed for purposes other than those listed in patient leaflets. If you have questions or concerns, or want more information about Carboplatin Injection, your physician and pharmacist have the complete prescribing information upon which this information is based. You may want to read it and discuss it with your doctor. Remember, no written summary can replace careful discussion with your your giver for

has been reported in patients recein cancer chemotherapy with mult drugs. It is not always possible to whether such effects are caused by ca-platin, another drug you may be tak-or your illness. Because some of the effects may be serious, you will n close medical supervision during to ment with Carboplatin Injection. rultip s to t f thes treat

Carbo-

- The most serious side effects of Caplatin Injection are:
 bleeding and reduced blood including reduced red blood (anemia) and platelets (neede proper blood clotting), which be severe enough to require t transfusion. You should tell your or right away if you notice any un bruising or bleeding, including tarry stools or blood in the urine infection Carboplatin Injectio temporarily lower the number of blood cells in your blood, increated during and after treatment the cor nurse will observe you carefu signs of allergic reaction;
 life-threatening allergic reaction;
 loss of hearing or ringing in the Contact your doctor right away if you experience any of these effects, or notice effects that v you or are troublesome.
 Of the less serious side effects at ated with Carboplatin Injection ment, the most common are na vomiting, diarrhea, loss of appetitic loss and numbness, tingling, burni pain in the hands or feet. A blood (d blood (s (needec), which r equire blc any unusu- uning blac ectir
 j black
 - tion can of white creasing

 - reaction t the doctor carefully for

 - in the it awa these that v ×0 ears

us side effects ass oplatin Injection tr common are naus a, loss of appetite, ass, tingling, burning or feet 3 associ-on treat-nausea, stite, hair stite, or